

# Famvir(R) for genital herpes

*famciclovir*

---

## Consumer Medicine Information

---

### What is in this leaflet

---

This leaflet answers some common questions about Famvir.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Famvir against the benefits it can provide.

**If you have any concerns about this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

---

### What Famvir is used for

---

Famvir is an antiviral medicine that is used to treat recurrent episodes of genital herpes in adults and adolescents. Genital herpes is a viral infection caused by herpes simplex 1 or 2. It is usually transmitted through sexual contact. Symptoms include tingling, burning or itching of the genitals, followed by blisters that may be painful. People who have frequent episodes of genital herpes can also take Famvir to help prevent the attacks.

Although Famvir does not cure the viral infection, it helps to relieve the symptoms and shorten their duration.

The best results are obtained if the medicine is started as soon as possible after the first symptoms begin to appear.

Taking Famvir does not prevent you from spreading the herpes virus to another person.

**Ask your doctor if you have any questions about why this medicine has been prescribed for you.**

Your doctor may have prescribed it for another reason.

Famvir is only available with a doctor's prescription. It is not addictive.

This medicine is not recommended for use in children under 12 years of age.

---

### Before you take Famvir

---

#### ***When you must not take it***

**Do not take Famvir if you have an allergy to:**

- \* famciclovir, the active ingredient, or to penciclovir, a related antiviral medicine
- \* any of the other ingredients of Famvir listed at the end of this leaflet

Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

**Do not take Famvir after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.**

In that case, return it to your pharmacist.

#### ***Before you start to take it***

**Tell your doctor if you have a problem with:**

- \* your body's immune system, which helps to fight off infections
- \* your kidneys
- \* your liver

Your doctor may want to take extra precautions in that case.

**Tell your doctor if you are pregnant, intend to become pregnant or you are breast-feeding.**

It is not known whether taking this medicine during pregnancy could affect your baby or whether the active ingredient in Famvir passes into the breast milk. Your doctor can discuss with you the benefits and risks involved.

**Tell your doctor if you are lactose intolerant.**

This medicine contains lactose.

**Tell your doctor if you are allergic to any other medicines, foods, dyes or preservatives.**

Your doctor will want to know if you are prone to allergies.

## ***Taking other medicines***

**Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.**

Some medicines and Famvir may interfere with each other. These include medicines that can affect your kidneys, such as probenecid.

You may need to take different amounts of these medicines or you may need to take different medicines. Your doctor and pharmacist have more information.

**If you have not told your doctor about any of these things, tell him/her before you start taking this medicine.**

---

## **How to take Famvir**

---

**Follow all directions given to you by your doctor and pharmacist carefully.**

These instructions may differ from the information contained in this leaflet.

**If you do not understand the instructions on the label, ask your doctor or pharmacist for help.**

### ***How much to take***

To treat an episode of genital herpes, the usual dose is one 125 mg tablet twice each day for five days, beginning as soon as possible (preferably within 6 hours) after the first symptoms appear.

To prevent recurrent episodes of genital herpes, the usual dose is one 250 mg tablet twice each day for up to 12 months.

For people whose immune system does not work as well as it should, the dose and duration of treatment may be increased.

For people who have reduced kidney function, the dose may be reduced.

### ***How to take it***

Unless your doctor tells you otherwise, take one tablet when you get up in the morning and one just before going to bed at night. Try to take the tablets at about the same time each day.

**Swallow the tablets whole with a full glass of water. You can take them with or without food.**

### ***How long to take it***

**Continue taking Famvir every day for as long as your doctor prescribes.**

To help clear up your infection, you must keep taking this medicine for the full course of treatment, even if your symptoms begin to clear up after a few days.

To prevent recurrent episodes of genital herpes, you must take the tablets each day, even if you have no symptoms.

### ***If you forget to take it***

**Take a dose as soon as you remember. Take your next tablet at the usual time, and then go back to taking it as you would normally.**

**Do not take a double dose to make up for the one that you missed.**

This may increase the chance of you getting an unwanted side effect.

**If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.**

### ***If you take too much (Overdose)***

**Immediately telephone your doctor or the Poisons Information Centre (telephone number 13 11 26), or go to Accident and Emergency at your nearest hospital if you think that you or anyone else may have taken too much Famvir. Do this even if there are no signs of discomfort or poisoning. Keep the telephone numbers for these places handy.**

---

## **While you are taking Famvir**

---

### ***Things you must do***

**If you become pregnant while taking Famvir, tell your doctor.**

Your doctor can discuss with you the risks of taking it while you are pregnant.

**If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking Famvir.**

**Tell any other doctor, dentist or pharmacist who treats you that you are taking Famvir.**

### ***Things you must not do***

**Do not give this medicine to anyone else even if their condition seems to be the same as yours.**

**Do not use it to treat any other complaints unless your doctor tells you to.**

### ***Things to be careful of***

**Be careful driving, operating machinery or doing jobs that require you to be alert until you know how Famvir affects you.**

This medicine can cause dizziness, sleepiness or confusion in some people.

**Avoid sexual intercourse when symptoms are present, even if you have started taking Famvir, because the risk of infecting your partner is increased. Use condoms between episodes to reduce the risk of infecting your partner.**

**Keep the areas affected by the virus as clean and dry as possible. Wear loose-fitting clothing to avoid irritating the blisters.**

**Avoid touching or scratching the sore area.**

You may spread the virus on your fingers.

---

## Side effects

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Famvir.**

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

**Do not be alarmed by these lists of possible side effects. You may not experience any of them. Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor if you notice any of the following and they worry you:**

- \* headache
- \* dizziness
- \* nausea (feeling sick) or vomiting
- \* diarrhoea

The above side effects are usually mild.

**Tell your doctor as soon as possible if you notice any of the following:**

- \* a rash on other parts of your body
- \* extreme sleepiness or confusion, usually in older people
- \* hallucinations (seeing or hearing things that are not really there)
- \* signs of a possible liver problem such as persistent pain in the upper right abdomen, yellowing of the skin and/or eyes, dark urine or pale bowel motions

The above side effects may need medical attention.

**Tell your doctor if you notice anything else that is making you feel unwell.**

Other side effects not listed here or not yet known may happen in some people.

---

## After using Famvir

---

### Storage

- \* Keep your medicine in the original container until it is time to take it.
- \* Store it in a dry place at room temperature.
- \* Do not store Famvir or any other medicine in the bathroom or near a sink.
- \* Do not leave it in the car or on window sills.

**Keep the medicine where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### Disposal

**If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine you have left over.**

---

## Product description

---

### What it looks like

Famvir 125 mg tablets are white round tablets, marked with either "Famvir" or "FV" on one side and "125" on the other; packs of 40 tablets.

Famvir 250 mg tablets are white round tablets, marked with either "Famvir" or "FV" on one side and "250" on the other; packs of 56 tablets.

### Ingredients

Famvir tablets contain 125 mg or 250 mg famciclovir as the active ingredient. They also contain:

- \* hydroxypropylcellulose
- \* lactose
- \* sodium starch glycollate

- \* magnesium stearate
- \* titanium dioxide
- \* hypromellose
- \* macrogol

Famvir tablets do not contain sucrose, gluten, tartrazine or any other azo dyes.

### More information

For more information about genital herpes, visit [www.herpes.com.au](http://www.herpes.com.au).

### Sponsor

Famvir is supplied in Australia by:

NOVARTIS Pharmaceuticals  
Australia Pty Limited

ACN 004 244 160

54 Waterloo Road

North Ryde NSW 2113

Telephone 1 800 671 203

Web site: [www.novartis.com.au](http://www.novartis.com.au)

(R) = Registered Trademark

This leaflet was prepared in  
August 2003

[fam(genital)290803c.doc] based on  
PI (fam290803i.doc)

Australian Registration Number.

Famvir 125 mg AUST R 54668

Famvir 250 mg AUST R 513891